

Inclusion-Exclusion Formulas from Independent Complexes

Dominique Attali^{*}
LIS laboratory
Domaine Universitaire, BP 46
38402 Saint Martin d'Hères, France
Dominique.Attali@lis.inpg.fr

Herbert Edelsbrunner[†]
Department of Computer Science
Duke University, Durham
Raindrop Geomagic, RTP, NC, USA
edels@cs.duke.edu

ABSTRACT

Using inclusion-exclusion, we can write the indicator function of a union of finitely many balls as an alternating sum of indicator functions of common intersections of balls. We exhibit abstract simplicial complexes that correspond to minimal inclusion-exclusion formulas. They include the dual complex, as defined in [2], and are characterized by the independence of their simplices and by geometric realizations with the same underlying space as the dual complex.

Categories and Subject Descriptors

F.2.2 [Analysis of Algorithms and Problem Complexity]: Non-numerical Algorithms and Problems—*Geometrical problems and computations, Computations on discrete structures*; G.2.1 [Discrete Mathematics]: Combinatorics—*Counting problems*

General Terms

Theory, Algorithms

Keywords

Combinatorial topology, discrete geometry, dual complexes, balls, spheres, indicator functions

1. INTRODUCTION

In this paper, we study inclusion-exclusion formulas for unions of finitely many balls in \mathbb{R}^d , generalizing previous results that derive such formulas from Delaunay triangulations and dual complexes.

Motivation. It is common in biochemistry to identify a molecule with the portion of space it occupies. This portion is sometimes

^{*}Partially supported by the IST Program of the EU under Contract IST-2002-506766 (Aim@Shape)

[†]Partially supported by NSF under grant CCR-00-86013 (Bio-Geometry).

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

SoCG '05, June 6–8, 2005, Pisa, Italy

Copyright 2005 ACM 1-58113-991-8/05/0006 ...\$5.00.

referred to as the *space-filling diagram*, and its simplest and most common form is a union of balls in \mathbb{R}^3 , each ball representing an atom of the molecule. The volume and surface area of this union are fundamental concepts that relate to physical forces acting on the molecules. We refer to [3] for a recent survey that describes this connection and also discusses derivatives of the volume and surface area, which are needed in simulating the motion of molecules.

Consider a finite set of balls in \mathbb{R}^3 and let us focus on the volume of the union. Generally, there are many inclusion-exclusion formulas that give the correct volume, even if we limit our attention to minimal formulas. The starting point of the work reported in this paper is the idea that this ambiguity could be useful in maintaining a formula for a moving set of balls. If we understand how long a formula remains valid, we can save time by delaying any changes until they become necessary. As a first step towards such an understanding, we study the family of minimal inclusion-exclusion formulas for a given set of balls.

Prior work. The principle of inclusion-exclusion is perhaps the most natural approach to measuring a union of balls. Letting B be a finite set of balls, the volume of the union is the alternating sum of volumes of intersections:

$$\text{vol} \bigcup B = \sum_{\emptyset \neq X \subseteq B} (-1)^{\text{card } X - 1} \text{vol} \bigcap X. \quad (1)$$

Writing n for the number of balls in B , we have $2^n - 1$ terms, each plus or minus the volume of the intersection of at most n balls. It seems the formula is only useful if all sets X with non-empty common intersection are small. More than a quarter century ago, Kratky [6] pointed out that even if this is not the case, one can substitute lower-order for higher-order terms and thus reduce the complexity of the formula. The software of Scheraga and collaborators [8] is based on this observation, but it is sometimes difficult to do the reduction correctly. In 1992, Naiman and Wynn [7] proved that Equation (1) is correct even if we limit the sum to sets X that correspond to simplices in the weighted Delaunay triangulation of B . By definition, this is the dual of the weighted Voronoi diagram of the balls, also known as the power diagram and the Dirichlet tessellation [1]. In the geometry literature, this dual is also known as the regular triangulation and the coherent triangulation of $\bigcup B$ [4]. In agreement with Kratky, this result implies that in \mathbb{R}^3 we only need sets X of cardinality at most four. Taking all such sets would lead to an incorrect formula, and Naiman and Wynn's result is a recipe for selecting sets that give a correct formula. In 1995, Edelsbrunner [2] further reduced the formula by proving that Equation (1) remains correct if we limit the sum to simplices in the dual complex, which

is a subcomplex of the weighted Delaunay triangulation of B . Besides giving a shorter formula, the terms obtained from the dual complex consist of balls that intersect in a unique pattern, which allows a simple implementation without case analysis [5].

Results. We refer to the specific intersection pattern exhibited by the balls in the dual complex formula as independent, a term whose technical definition will be given shortly. Our first result generalizes this formula to a family of formulas whose terms exhibit the same pattern. Specifically, *if K is an abstract simplicial complex whose simplices are independent sets of balls and whose canonical geometric realization has the same boundary complex and underlying space as the dual complex then the corresponding inclusion-exclusion formula is correct.* To prove that this is a proper generalization, we show in Figure 1 that even already for four disks in the plane we can have more than one such formula. Our second result states that *the inclusion-exclusion formulas in the family specified in our first result are minimal and exhaust all minimal formulas that correspond to simplicial complexes.*




Figure 1: Four disks that permit two correct, minimal inclusion-exclusion formulas. The upper complex on the right is the dual complex of the disks and corresponds to the formula $a + b + c + d - ab - ad - bc - bd - cd + abd + bcd$, in which we write a for the area of disk a , ab for the area of the intersection of a and b , etc. To get the formula of the lower complex, we substitute $-ac + abc + acd$ for $-bd + abd + bcd$.

Outline. Section 2 presents definitions and the formal statements of our two results. Section 3 proves the first result and Section 4 proves the second. Section 5 concludes this paper.

2. STATEMENT OF RESULTS

In this paper, a simplex may be abstract (a collection of balls) or geometric (the convex hull of affinely independent points). We use both interpretations interchangeably and introduce notation that does not distinguish between them.

Independent simplices. Let B be a finite set of closed balls in \mathbb{R}^d . Throughout this paper, we assume that the balls are in general position, which includes that every ball has positive radius and the common intersection of any $k + 1$ bounding spheres is either empty or a sphere of dimension $d - k - 1$. In particular, this prevents the common intersection degenerating to a single point. An *abstract simplex* is a non-empty subset $\alpha \subseteq B$ and its *dimension* is one less than its cardinality, $\dim \alpha = \text{card } \alpha - 1$. A k -*simplex* is an abstract simplex of dimension k . It is *independent* if for every subset $\gamma \subseteq \alpha$, including $\gamma = \emptyset$, there is a point that belongs to all balls in γ but not to any ball not in γ :

$$\bigcap \gamma - \bigcup (\alpha - \gamma) \neq \emptyset.$$

By assumption of general position, we have an open set of points for each $\gamma \subseteq \alpha$. In \mathbb{R}^2 , there are only three types of independent simplices, one each for one, two, and three disks, as shown in Figure 2. Four disks cannot be independent because the four bounding




Figure 2: From left to right: an independent simplex of dimension $k = 0, 1, 2$.

circles decompose the plane into at most 14 regions but we need 16, one each for the 2^4 subsets. Similarly, in \mathbb{R}^d we have $d + 1$ types of independent simplices. For $0 \leq k \leq d - 1$, we can construct an independent k -simplex in \mathbb{R}^d from one in \mathbb{R}^k , by drawing the latter in a k -dimensional plane and replacing each k -dimensional ball by a d -dimensional ball, using the same center and radius. For $k = d$, we get an independent d -simplex by adding a single ball to an independent $(d - 1)$ -simplex. The $d(d - 1)$ -spheres bounding the balls of the $(d - 1)$ -simplex intersect in two points, and the extra ball contains one of the two points in its interior and does not contain the other point. If $\beta \subseteq \alpha$ is a non-empty subset, we call β a *face* of α and α a *coface* of β . Clearly, every face of an independent simplex is independent.

General position. We think of \mathbb{R}^d as the subspace of \mathbb{R}^{d+1} spanned by the first d coordinate axes. For each d -dimensional ball b_i with center $z_i \in \mathbb{R}^d$ and radius $r_i > 0$, we construct the corresponding *ghost sphere*,

$$s_i = \{y \in \mathbb{R}^{d+1} \mid \|y - z_i\|^2 - r_i^2 = 0\},$$

which is a d -sphere in \mathbb{R}^{d+1} . Using this concept, we can now define what exactly we mean by a set of balls to be in *general position*, namely that the common intersection of any $k + 1$ ghost spheres is either empty or a sphere of dimension $d - k$. We note that for $0 \leq k < d$, this is equivalent to the condition we mentioned earlier: all radii are positive and the common intersection of any $k + 1$ bounding spheres is either empty or a sphere of dimension $d - k - 1$. For $k \geq d$ we get new conditions. We need some definitions before we can explain them in terms of concepts intrinsic to \mathbb{R}^d .

Call $\pi_i(x) = \|x - z_i\|^2 - r_i^2$ the *power distance* of the point $x \in \mathbb{R}^d$ from b_i and note that $b_i = \pi_i^{-1}(-\infty, 0]$ and the zero-set of π_i is the $(d - 1)$ -sphere bounding b_i . Using the power distance, we decompose $\bigcup B$ into convex cells, one for each ball. Specifically, the cell of b_i consists of all points $x \in b_i$ with $\pi_i(x) \leq \pi_j(x)$ for all $b_j \in B$. It is not difficult to see that the cell of b_i is the intersection of b_i with a convex polyhedron, keeping in mind that this polyhedron or its intersection with b_i may be empty. To describe the relation between this cell and the ghost sphere of b_i , we define $\sigma_i(y) = \|y - x\|^2 + \pi_i(x)$, where x is the orthogonal projection of $y \in \mathbb{R}^{d+1}$ to \mathbb{R}^d . The ghost sphere itself is the zero-set, $s_i = \sigma_i^{-1}(0)$. For a given point $x \in b_i$, define $y_i(x) \in \mathbb{R}^{d+1}$ above \mathbb{R}^d such that $\|y_i(x) - x\|^2 + \pi_i(x) = 0$; it is the point on the upper hemi-sphere of s_i whose orthogonal projection to \mathbb{R}^d is x . The condition for x to belong to the cell of b_i now translates to $\|y_i(x) - x\|^2 \geq \|y_j(x) - x\|^2$ whenever $y_j(x)$ is defined. In words, the cell of b_i is the orthogonal projection of s_i 's contribution to the upper envelope of the ghost spheres, as illustrated in Figure 3.




Figure 3: Upper envelope of the ghost spheres of three independent disks and the corresponding decomposition of the union of disks into convex cells.

Let us return to the case $k \geq d$ of our general position assumption. It says that the common intersection of any $k + 1 \geq d + 1$ ghost spheres is either empty or a 0-sphere, and the latter case can only happen if $k = d$. Equivalently, the common intersection of the cells decomposing the union of any $k + 1 \geq d + 1$ balls is either empty or a point in the interior of the union, and the latter case can happen only if $k = d$.

Characterizing independence. Besides for expressing our general position assumption, ghost spheres can be used for characterizing independent simplices. This characterization will be important in establishing the Non-nesting Lemma in Section 3, a crucial step in the proof of our first result.

GHOST SPHERE LEMMA. A k -simplex of $k + 1$ balls in general position is independent iff the common intersection of its $k + 1$ ghost spheres is a sphere of dimension $d - k$.

PROOF. All ghost spheres have their centers in \mathbb{R}^d , which implies that the arrangement of the $k + 1$ d -spheres is symmetric with respect to \mathbb{R}^d . The number of chambers (cells of dimension $d + 1$) in this arrangement is the same above and below \mathbb{R}^d , and indeed the same altogether. To prove the claim, we show that there are 2^{k+1} chambers iff the $k + 1$ ghost spheres meet in a $(d - k)$ -sphere.

We prove one direction by counting the chambers while adding one d -sphere at a time. Letting s_0, s_1, \dots, s_k be this sequence, we note that s_0 creates two chambers, one inside and one outside. When we add s_j , we consider its decomposition into d -dimensional patches defined by the preceding d -spheres. As we add the patches, again one by one, each patch may or may not cut a chamber into two. To reach the necessary 2^{k+1} chambers, we need to double the number of chambers each time we add a d -sphere. This is only possible if s_j is decomposed into 2^j patches, the maximum possible, and each patch cuts a chamber into two. Using stereographic projection, we map s_j to a d -dimensional plane and its patches to the $(d$ -dimensional) chambers in the arrangement of j $(d - 1)$ -spheres, the images of the $s_j \cap s_i$ for $0 \leq i \leq j - 1$. By induction over the dimension, having 2^j such chambers implies the j $(d - 1)$ -spheres meet in a common $(d - j)$ -sphere. In the last step, we have $j = k$ and get a $(d - k)$ -sphere common to all $k + 1$ d -spheres.

Proving the reverse implication is easier. If $k = d$, we have $d + 1$ d -spheres meeting in a 0-sphere, that is, a pair of points. In a sufficiently small neighborhood of one of these two points, the d -spheres behave like d -dimensional planes, decomposing the neighborhood into 2^{d+1} orthants. Each orthant corresponds to a unique subset of the d -spheres and belongs to a unique chamber in the arrangement they define. It follows that the corresponding d -simplex is independent. If $k < d$, we pick a point on the common

$(d - k)$ -sphere and intersect the arrangement with the $(k + 1)$ -dimensional plane that passes through this point and the centers of the $k + 1$ d -spheres. Within this plane, we have $k + 1$ k -spheres meeting in a common 0-sphere and we apply the above argument to conclude that the corresponding k -simplex is independent. \square

Simplicial complexes. An *abstract simplicial complex* is a collection of non-empty abstract simplices, K , that contains, with every simplex, the faces of that simplex. If B is the set of vertices then K is a subset of the power set, $K \subseteq 2^B$. Figure 4 illustrates the definitions. A *geometric realization* maps every abstract simplex to




Figure 4: A finite set of disks and the canonical realization of an abstract simplicial complex over that set. The vertices, edges, and triangles correspond to abstract simplices of dimension 0, 1, and 2. Take a moment to verify that all simplices are independent.

a geometric simplex of the same dimension such that the intersection of the images of two abstract simplices α and β is the image of $\alpha \cap \beta$, which is either empty or a face of both. In this paper, the vertices are closed balls and we map every abstract simplex to its *canonical image*, defined as the convex hull of the centers of its balls. We call K *canonically realizable* if this map is a geometric realization. We use the letters α and β to denote the abstract simplices as well as their images, which are geometric simplices. Similarly, we use the letter K to denote the abstract simplicial complex as well as its geometric realization, which is a geometric simplicial complex. Its *underlying space* is the set of points covered by the geometric simplices, which we denote as $|K|$. The *star* of an abstract simplex β is the set of cofaces $\alpha \in K$, and the *link* of β is the set of simplices $\alpha - \beta$ with $\alpha \supset \beta$. Assuming K is geometrically realized in \mathbb{R}^d , the link of every k -simplex is a triangulation of the sphere of dimension $d - k - 1$ or a proper subcomplex of such a triangulation. We define the *boundary complex* of K as the subset of simplices in the latter category. This is also the subcomplex of simplices contained in the boundary of $|K|$.

Dual and other independent complexes. Let B be a set of closed balls and recall the decomposition of $\bigcup B$ into convex cells described above. The nerve of this collection of cells is particularly important for the developments reported in this paper. The *dual complex* of B is the canonical realization of this nerve, obtained by mapping every $k + 1$ cells with non-empty intersection to the k -simplex spanned by the centers of the corresponding balls. This construction is illustrated in Figure 5, where we see the dual complex superimposed on the decomposition of the union into convex

cells. It is perhaps not obvious but true that the canonical mapping of abstract simplices defines a geometric realization of the nerve, provided the balls in B are in general position.




Figure 5: The dual complex of the disks in Figure 4. Its simplices record the overlap pattern of the cells in the decomposition of the union. In this example, the dual complex has the same boundary complex and underlying space as the independent complex in Figure 4 but differs from it in six edges and twelve triangles.

Given a finite set of balls in general position, B , we are primarily interested in abstract simplicial complexes K of B that satisfy the following three conditions:

Independence: all simplices in K are independent;

Realizability: K is canonically realizable in \mathbb{R}^d ;

Boundary: the boundary complex and underlying space of K are the same as those of the dual complex.

An *independent complex* is an abstract simplicial complex that satisfies the independence condition. We note that there is an alternative way to express the boundary condition, without references to the dual complex, by comparing the boundaries of K and $\bigcup B$. In particular, a simplex α belongs to the boundary complex of K iff there is a point on the boundary of $\bigcup B$ that belongs to all balls in α and to no others.

First result: correct indication. The *indicator function* of a subset $A \subseteq \mathbb{R}^d$ is the map $\mathbf{1}_A : \mathbb{R}^d \rightarrow \{0, 1\}$ defined by

$$\mathbf{1}_A(x) = \begin{cases} 1 & \text{if } x \in A, \\ 0 & \text{if } x \notin A. \end{cases}$$

Given a finite set of balls in \mathbb{R}^d , our first result states that the inclusion-exclusion formula defined by a simplicial complex that satisfies the above three conditions gives the correct indicator function of the union.

THEOREM A. Let B be a finite set of closed balls in general position in \mathbb{R}^d and K an independent complex that is canonically realizable in \mathbb{R}^d and satisfies the boundary condition. Then $\mathbf{1}_{\bigcup B} = \sum_{\alpha \in K} (-1)^{\dim \alpha} \mathbf{1}_{\cap \alpha}$.

Using Theorem A, we obtain formulas for the volume or other mea-

asures of the union by integrating the density function, $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\begin{aligned} \text{meas} \bigcup B &= \int_{x \in \bigcup B} \mu(x) \, dx \\ &= \int_{x \in \mathbb{R}^d} \mu(x) \mathbf{1}_{\bigcup B}(x) \, dx \\ &= \sum_{\alpha \in K} (-1)^{\dim \alpha} \int_{x \in \mathbb{R}^d} \mu(x) \mathbf{1}_{\cap \alpha}(x) \, dx \\ &= \sum_{\alpha \in K} (-1)^{\dim \alpha} \text{meas} \cap \alpha. \end{aligned}$$

For $d = 2$, the edge skeleton of K is a planar graph implying that the number of terms in the above formula is less than six times $n = \text{card } B$. More generally, the number of terms is bounded from above by some constant times $n^{\lceil d/2 \rceil}$.

Second result: minimality. The inclusion-exclusion formula that corresponds to an abstract simplicial complex $K \subseteq 2^B$ gives a map $\text{IEF}_K : \mathbb{R}^d \rightarrow \mathbb{Z}$ defined by

$$\text{IEF}_K(x) = \sum_{\alpha \in K} (-1)^{\dim \alpha} \mathbf{1}_{\cap \alpha}(x).$$

The formula is *minimal* if $\text{IEF}_L \neq \text{IEF}_K$ for all proper subsets $L \subset K$. By Theorem A, we have $\text{IEF}_K = \mathbf{1}_{\bigcup B}$ if K is an independent complex canonically realizable in \mathbb{R}^d that satisfies the boundary condition. Our second result states that such complexes have minimal formulas and that they exhaust the family of complexes with minimal formulas.

THEOREM B. Let B be a finite set of closed balls in general position in \mathbb{R}^d and $K \subseteq 2^B$ an abstract simplicial complex with $\text{IEF}_K = \mathbf{1}_{\bigcup B}$. This formula is minimal iff K is independent, canonically realizable in \mathbb{R}^d , and satisfies the boundary condition.

3. PROOF OF THEOREM A

In this section, we present our proof of Theorem A. Starting with a finite set of balls, we first add small balls covering the rest of \mathbb{R}^d to get an infinite but discrete set, and we second use this discrete set as the basis for a continuous set. Both steps are instrumental in obtaining the technical results that imply Theorem A.

Induced subcomplexes. Given an abstract simplicial complex $K \subseteq 2^B$, a subset $B_0 \subseteq B$ induces the subcomplex $K_0 = K \cap 2^{B_0}$. To establish our first result, we associate to each point $x \in \mathbb{R}^d$ the subset $B_x \subseteq B$ of balls that contain x and the subcomplex $K_x \subseteq K$ induced by B_x . We have

$$\begin{aligned} \text{IEF}_K(x) &= \sum_{\alpha \in K} (-1)^{\dim \alpha} \mathbf{1}_{\cap \alpha}(x) \\ &= \sum_{\alpha \in K_x} (-1)^{\dim \alpha}. \end{aligned}$$

The latter sum is the Euler characteristic of K_x , which we denote as $\chi(K_x)$. For all points $x \notin \bigcup B$, $K_x = \emptyset$ and $\text{IEF}_K(x) = \chi(K_x) = 0$. To tackle the points inside the union, we recall that the Euler characteristic of every contractible set is 1. As explained later, such a set has the homotopy type of a point, which in the plane includes trees and closed disks. We will show that $|K_x|$ is contractible, for every point $x \in \bigcup B$, which will then imply Theorem A.

The union of the balls in B_x is star-convex, which implies that $\bigcup B_x$ is contractible. In spite of the fact that the underlying space

of the subcomplex K_x induced by B_x is not necessarily star-convex, we will be able to prove that $|K_x|$ is also contractible. Before embarking on this proof, we introduce the discrete and continuous sets of balls. Using the continuous set, we will find a set between $|K_x|$ and $\bigcup B_x$, which we will show is star-convex and of the same homotopy type as $|K_x|$.

From finite to discrete sets of balls. A simplicial complex is *locally finite* if the star of every vertex is finite. We extend the finite set of balls B to a discrete set $\bar{B} = B \cup B_\varepsilon$. Simultaneously, we construct a locally finite independent complex $\bar{K} \supseteq K$ whose vertices are the balls in \bar{B} and whose underlying space is \mathbb{R}^d . The construction depends on a positive number ε , the radius of the balls in B_ε added to B . We require that \bar{B} covers \mathbb{R}^d while the center of every ball in B_ε lies outside all other balls in \bar{B} , as illustrated in Figure 6. Choosing $\varepsilon > 0$ sufficiently small, we construct B_ε one




Figure 6: Extension of the independent complex in Figure 4 by adding disks of radius ε . The rectangular frame delimits the portion of the configuration reused in Figure 7.

ball at a time, picking the center outside all previous balls, until \bar{B} covers \mathbb{R}^d . Assuming the balls in B are in general position, it is clear that we can construct \bar{B} such that its balls are also in general position. To see what ε is sufficiently small, we consider the cells in the decomposition of $\bigcup B$. As we add balls of radius ε , these cells give up territory to the new balls, but not more than what is covered by the new balls. By shrinking ε , we can make the loss of territory as narrow as we like. By assumption of general position, we can therefore maintain the non-empty common intersection of any collection of cells in the decomposition of $\bigcup B$. It follows that the dual complex of B is a subcomplex of the dual complex of \bar{B} . The boundary complex of the dual complex of B is the same as that of K . We can therefore construct \bar{K} equal to K inside and equal to the dual complex of \bar{B} outside that boundary, as illustrated in Figure 6. We finally note the choice of balls implies that \bar{K} is locally finite.

From discrete to continuous sets of balls. An abstract simplex is a finite set of balls, $\alpha = \{b_0, b_1, \dots, b_k\}$. We extend α to an infinite set by considering convex combinations of balls in α . Recall that $\pi_i : \mathbb{R}^d \rightarrow \mathbb{R}$ maps each point $x \in \mathbb{R}^d$ to the power distance from b_i and that $b_i = \pi_i^{-1}(-\infty, 0]$. An *affine combination* of α is a ball $b = \pi^{-1}(-\infty, 0]$ for which there are real numbers λ_i , summing to 1, such that $\pi = \sum_{i=0}^k \lambda_i \pi_i$. Recall also that the ghost sphere s_i of b_i is the zero-set of the map $\sigma_i : \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ defined by $\sigma_i(y) = \|y - x\|^2 + \pi_i(x)$, where x is the orthogonal projection of y onto \mathbb{R}^d . Similarly, the ghost sphere s of b is the zero-set

of $\sigma = \sum_{i=0}^k \lambda_i \sigma_i$. A point y belongs to the common intersection of the s_i iff $\sigma_i(y) = 0$, for all i , which implies

$$\bigcap_{i=0}^k s_i \subseteq s. \quad (2)$$

A *convex combination* b of α is an affine combination for which all λ_i are non-negative. If a point x belongs to all balls in α then $\pi_i(x) \leq 0$, for all i , which implies $\pi(x) \leq 0$. Furthermore, if $\pi(x) \leq 0$ then $\pi_i(x) \leq 0$ for at least one index i , which implies that x belongs to at least one ball in α . In set notation,

$$\bigcap \alpha \subseteq b \subseteq \bigcup \alpha. \quad (3)$$

Letting $\alpha \in \bar{K}$ be the simplex whose interior contains the point $z \in \mathbb{R}^d$, we write b_z for the (unique) convex combination of balls in α whose center is z . Incidentally, the coefficients that define b_z in terms of the balls in α are the same as the ones that define z in terms of the centers of the balls in α . Relation (2) is useful when we consider a line and the balls b_z whose centers z lie on the line. These balls intersect the line in intervals. It turns out that as we move the center monotonically along the line, the left endpoint also moves monotonically and so does the right endpoint. It is convenient to prove this for the extension \bar{K} of K for which there are balls for all points along the line. As usual, we assume the balls in \bar{B} are in general position.

NON-NESTING LEMMA. For any two points $x \neq y$ in \mathbb{R}^d , the two balls b_x and b_y are either disjoint or independent.

PROOF. Consider first the case in which x and y are points of a common d -simplex α in \bar{K} . Since α is independent, the ghost spheres of α intersect in a common 0-sphere. By Relation (2), the ghost spheres of b_x and b_y pass through this 0-sphere and thus meet in a $(d-1)$ -sphere. It follows that b_x and b_y are independent.

If x and y do not belong to a common d -simplex, there is a point z on the line segment connecting x and y that lies on a $(d-1)$ -simplex. The number of $(d-1)$ -simplices separating x from z is strictly smaller than the number separating x from y , and similar for z and y . We can therefore use induction to show that b_x and b_z as well as b_z and b_y are either disjoint or independent. Since z lies between x and y , it follows that also b_x and b_y are either disjoint or independent. \square

Intermediate set of centers. For a point $x \in \mathbb{R}^d$, we write $\bar{B}_x \subseteq \bar{B}$ for the set of balls that contain x and $\bar{K}_x \subseteq \bar{K}$ for the subcomplex induced by \bar{B}_x . We prove that $|\bar{K}_x|$ is contractible by showing it has the same homotopy type as

$$\bar{Z}_x = \{z \in \mathbb{R}^d \mid x \in b_z\},$$

which we later prove is star-convex. Let α be the simplex whose interior contains the point $z \in \mathbb{R}^d$. By Relation (3), $z \in \bar{Z}_x$ if all balls in α contain x . Similarly, $z \notin \bar{Z}_x$ if none of the balls in α contains x . As illustrated in Figure 7, the first property implies $|\bar{K}_x| \subseteq \bar{Z}_x$. Let \bar{L}_x be the subcomplex of \bar{K} induced by $\bar{B} - \bar{B}_x$. Each vertex of \bar{K} is either in \bar{K}_x or in \bar{L}_x . It follows that each abstract simplex α that is in neither induced subcomplex is the union of its largest faces $\alpha_K \in \bar{K}_x$ and $\alpha_L \in \bar{L}_x$. The corresponding geometric construction writes α as the union of line segments pq connecting points $p \in \alpha_K$ with points $q \in \alpha_L$. We note that b_p contains x and b_q does not contain x . The boundary spheres of b_p and b_q meet in a $(d-2)$ -sphere which is also contained in the bounding sphere of every ball along the line segment




Figure 7: The point x is contained in four disks, which induce the dark subcomplex \bar{K}_x , consisting of two triangles and their faces. The other disks induce the light subcomplex \bar{L}_x . We get \bar{Z}_x by adding initial portions of the line segments covering the in-between simplices to the underlying space of \bar{K}_x .

between p and q . As we move from p to q , the ball shrinks on the side of x , which implies that there is a unique point y on pq such that $x \in b_z$ for all z between p and y , including y , and $x \notin b_z$ for all z between y and q , excluding y . In other words, \bar{Z}_x can be written as $|\bar{K}_x|$ union all line segments py as described. These line segments can be shrunk continuously towards $|\bar{K}_x|$. Formally, we define $y(\lambda) = (1 - \lambda)y + \lambda p$, for $0 \leq \lambda \leq 1$, and we consider the set $\bar{Z}_x(\lambda)$ defined as $|\bar{K}_x|$ union all line segments $py(\lambda)$. We have $\bar{Z}_x(0) = \bar{Z}_x$ and $\bar{Z}_x(1) = |\bar{K}_x|$. We thus constructed a deformation retraction that takes \bar{Z}_x to $|\bar{K}_x|$, proving that the two have the same homotopy type.

HOMOTOPY TYPE LEMMA. $|\bar{K}_x| \simeq \bar{Z}_x$.

As mentioned earlier, a contractible set has the homotopy type of a point. By the above lemma, $|\bar{K}_x|$ is contractible iff \bar{Z}_x is contractible. We prove the latter by showing that \bar{Z}_x is the union of line segments emanating from a common endpoint. This implies that \bar{Z}_x is contractible because we can again exhibit a deformation retraction by shrinking the line segments, this time toward their common endpoint.

STAR-CONVEXITY LEMMA. \bar{Z}_x is star-convex.

PROOF. Observing that $x \in \bar{Z}_x$, we show that any line that passes through x intersects \bar{Z}_x in a single line segment. To reach a contradiction, assume there are points y and z on such a line through x such that z lies strictly between x and y and $x \in b_y$ but $x \notin b_z$. Then $b_z \subseteq b_y$, which contradicts the Non-nesting Lemma. \square

It is not too difficult to show that the boundary of \bar{Z}_x is piecewise linear, as suggested by Figure 7. In other words, \bar{Z}_x is a star-convex polytope.

Finale. We finally state and prove the crucial technical result that implies Theorem A.

CONTRACTIBILITY LEMMA. Let B be a finite set of closed balls in general position in \mathbb{R}^d . For every point $x \in \bigcup B$, the underlying space of the subcomplex K_x induced by the balls that contain x is contractible.

PROOF. We first establish the result for points x in the interior of $\bigcup B$. We may assume that $\varepsilon > 0$ is sufficiently small such that none of the balls in B_ε contains x . Hence, $\bar{B}_x = B_x$ and $\bar{K}_x = K_x$. By the Homotopy Type Lemma, $|K_x|$ and \bar{Z}_x have the same homotopy type, and by the Star-convexity Lemma, \bar{Z}_x is star-convex and therefore contractible. It follows that $|K_x|$ is contractible. By assumption of general position, every point x on the boundary has a point y in the interior of $\bigcup B$ that is contained in the same balls of B as x . Therefore, $K_x = K_y$ and the claim follows by the first argument. \square

As mentioned earlier, the contractibility of $|K_x|$ implies $\text{IEF}_K(x) = \chi(K_x) = 1$ for all points $x \in \bigcup B$. Theorem A follows.

4. PROOF OF THEOREM B

In this section, we present a proof of Theorem B. We begin by establishing Equation (4) as our main technical tool.

Witness points. Let B be a finite set of closed balls in general position in \mathbb{R}^d , as usual. Let $\beta \subseteq B$ be an independent k -simplex and $l(\beta)$ the $(d - k - 1)$ -sphere common to the $(d - 1)$ -spheres bounding the balls in β . By assumption of general position, almost all points of $l(\beta)$ do not lie on any other bounding $(d - 1)$ -sphere, and we let $y \in l(\beta)$ be one such point. We consider 2^{k+1} points x_γ near y , one for each subset $\gamma \subseteq \beta$, as illustrated in Figure 8. We require that the points witness the independence of β , that is, $x_\gamma \in \bigcap \gamma - \bigcup(\beta - \gamma)$ for all γ , and that every other ball in $B - \beta$ either contains all of the points or none of them. Given a collection of independent simplices $L \subseteq 2^B$, we study the alternating sum

$$\begin{aligned} \chi &= \sum_{\gamma \subseteq \beta} (-1)^{\dim \gamma} \text{IEF}_L(x_\gamma) \\ &= \sum_{\gamma \subseteq \beta} (-1)^{\dim \gamma} \sum_{\alpha \in L} (-1)^{\dim \alpha} \mathbf{1}_{\bigcap \alpha}(x_\gamma) \\ &= \sum_{\alpha \in L} (-1)^{\dim \alpha} \kappa(\alpha), \end{aligned}$$

where $\kappa(\alpha) = \sum_{\gamma \subseteq \beta} (-1)^{\dim \gamma} \mathbf{1}_{\bigcap \alpha}(x_\gamma)$ and $\dim \emptyset = -1$. If $\bigcap \alpha$ does not contain y then it contains none of the points x_γ and we have $\mathbf{1}_{\bigcap \alpha}(x_\gamma) = 0$ for all γ . Otherwise, there is a unique largest subset $\delta \subseteq \beta$ contained in α , namely $\delta = \beta \cap \alpha$, and we have $x_\gamma \in \bigcap \alpha$ iff $\delta \subseteq \gamma \subseteq \beta$. The set of such γ has the structure of an abstract simplex and we have $\kappa(\alpha) = 0$ unless $\delta = \beta$. Equivalently, $\kappa(\alpha) = 0$ unless $\beta \subseteq \alpha$ in which case $\kappa(\alpha) = (-1)^{\dim \beta} \mathbf{1}_{\bigcap \alpha}(x_\beta)$. Letting $L_\beta \subseteq L$ be the set of simplices α that contain β , we get

$$\chi = (-1)^{\dim \beta} \text{IEF}_{L_\beta}(y), \quad (4)$$

because x_β and y are contained in the same balls. We are interested in two special cases. The first case is characterized by IEF_L being constant in a neighborhood of y . By the choice of points x_γ , we have $\text{IEF}_L(x_\gamma) = \text{IEF}_L(y)$ for all γ . Plugging the common value into the definition, we get $\chi = 0$, and using Equation (4), we get $\text{IEF}_{L_\beta}(y) = 0$. We state this result in words, letting β be an independent simplex and $y \in l(\beta)$ a point not on the bounding sphere of any ball in $B - \beta$, as before.

EVEN COROLLARY. If IEF_L is constant in a neighborhood of y then the number of cofaces $\alpha \in L$ of β with $y \in \bigcap \alpha$ that have even dimension is the same as the number of such cofaces that have odd dimension.

The name of the claim is motivated by the weaker implication that the number of cofaces α of β with $y \in \bigcap \alpha$ is even. The second special case is characterized by $\text{IEF}_L(x_\gamma) = \text{IEF}_L(y)$ for all $\gamma \neq \emptyset$ and $\text{IEF}_L(x_\emptyset) = \text{IEF}_L(y) - 1$. This arises, for example, when the inclusion-exclusion formula of L is the indicator function of $\bigcup B$ and y lies on the boundary of the union. Plugging the values into the definition, we get $\chi = 1$, and using Equation (4), we get $\text{IEF}_{L_\beta}(y) = \pm 1$. We state a weaker implication in words.

ODD COROLLARY. If IEF_L is constant around y , except in the orthant of x_\emptyset where it is one less, then the number of cofaces $\alpha \in L$ of β with $y \in \bigcap \alpha$ is odd.

Redundant subsets. A subset L of an abstract simplicial complex K is *redundant* if $\text{IEF}_K = \text{IEF}_{K-L}$. Equivalently,

$$\text{IEF}_L = \sum_{\alpha \in L} (-1)^{\dim \alpha} \mathbf{1}_{\bigcap \alpha}$$

vanishes everywhere. We use the Even Corollary to derive structural properties of redundant subsets.

REDUNDANT SUBSET LEMMA. Let B be a finite set of closed balls in general position in \mathbb{R}^d , K an independent complex over B , and L a redundant subset of K .

- (i) If L contains a k -simplex β , with $k < d$, then L contains at least one proper coface $\alpha \supset \beta$.
- (ii) If L contains a $(d-1)$ -simplex β , then L contains two d -simplices whose canonical images in \mathbb{R}^d intersect in the canonical image of β .
- (iii) If L contains a d -simplex α , then L contains all $d+1$ $(d-1)$ -faces of α .

PROOF. To get (i), let $y \in l(\beta)$. Since IEF_L vanishes everywhere, and therefore also in a neighborhood of y , the Even Corollary implies that L contains an even number of cofaces α of β with $y \in \bigcap \alpha$. One such cofaces is β itself, which implies the number is at least two and therefore includes at least one proper coface.

To get (ii), observe that $l(\beta)$ consists of two points, y and z . Applying the above argument to y we obtain a d -simplex $\alpha \supset \beta$ in L . Since $d-1$ and d are the only dimensions to consider, and for trivial reasons β is the only $(d-1)$ -simplex that contains β , the d -simplex α is unique. Since α is independent, the extra ball in α contains y and does not contain z . Symmetrically, we get a unique d -simplex whose extra ball contains z and does not contain y . The centers of the two extra balls lie on opposite sides of the $(d-1)$ -dimensional plane spanned by β . It follows that the two d -simplices lie on opposite sides of the $(d-1)$ -simplex, as illustrated in Figure 8 on the right.

To get (iii), we consider the common intersection of the $d+1$ balls in α . Since α is independent, this intersection has the shape of a d -simplex with spherical faces. For each of its $d+1$ vertices y , we consider the $(d-1)$ -face β of α with $y \in l(\beta)$. The Even Corollary implies that L contains at least one coface of β , besides α , whose common intersection contains y . As proved above, α is the only proper coface of β with $y \in \bigcap \alpha$, leaving β itself as the only remaining possibility. \square

Sufficiency. We are ready to prove one direction of Theorem B. Specifically, we show that an abstract simplicial complex K that is independent, canonically realizable in \mathbb{R}^d , and satisfies the boundary condition has a minimal inclusion-exclusion formula. Equivalently, such a complex K contains no redundant subset.

To obtain a contradiction, we assume K has a non-empty redundant subset L . Because of (i) in the Redundant Subset Lemma, we may assume that L contains at least one d -simplex. Using (iii) of the same lemma, we see that L also contains the $(d-1)$ -faces of that d -simplex. By iterating (ii) and (iii), we conclude that L contains all d -simplices of a component formed by connecting the d -simplices across shared $(d-1)$ -faces. But then L also contains the boundary $(d-1)$ -simplices of that component, which exist because K is finite and geometrically realized in \mathbb{R}^d . But now we arrived at a contradiction because a boundary $(d-1)$ -simplex lacks the d -simplex on its other side which, by (ii) of the Redundant Subset Lemma, ought to be in L .

Boundary and interior. Having established one direction of Theorem B, we now prepare the other. Let B be a finite set of balls in \mathbb{R}^d and $K \subseteq 2^B$ an abstract simplicial complex. The only properties we assume are that the balls are in general position and that $\text{IEF}_K = \mathbf{1}_{\bigcup B}$.

INSIDE-OUTSIDE LEMMA. Let $\beta \subseteq B$ be an independent k -simplex, with $k < d$, not necessarily in K , and $y \in l(\beta)$ a point not on the $(d-1)$ -sphere bounding any ball in $B - \beta$.

- (i) If y lies in the interior of $\bigcup B$ then $\beta \in K$ implies that K contains a proper coface $\alpha \supset \beta$.
- (ii) If y lies on the boundary of $\bigcup B$ then $\beta \in K$.

PROOF. To get (i), we note that IEF_K is equal to 1 in a neighborhood of y . The Even Corollary implies that K contains an even number of cofaces of β whose common intersections contain y . If β is in K then this number is at least two so there is also a proper coface $\alpha \supset \beta$ in K . To get (ii), we note that IEF_K is equal to 1 in a neighborhood of y except outside $\bigcup B$, where it is 0. The Odd Corollary implies that K contains an odd number of cofaces of β whose common intersections contain y . This odd number is at least one, and since K is a complex, this implies that K also contains β . \square




Figure 8: The edge belongs to 0, 1, or 2 triangles depending on whether 2, 1, or 0 of the points y and z lie on the boundary of the union of disks. The four points near y are the points x_γ used in the derivation of Equation (4).

Similar to before, it is possible to get more detailed information when $\beta \in K$ is a $(d-1)$ -simplex. Then $l(\beta)$ consists of two points, y and z , and we get 0, 1, or 2 d -simplices sharing β depending on whether both points lie on the boundary, one lies on the boundary and the other in the interior, or both lie in the interior of $\bigcup B$. The three cases are illustrated in Figure 8.

Necessity. We are finally ready to prove the second direction of Theorem B. Specifically, we show that an abstract simplicial complex K with minimal inclusion-exclusion formula $\text{IEF}_K = \mathbf{1}_{\bigcup B}$

is independent, canonically realizable, and has the same boundary complex and underlying space as the dual complex.

First independence. Suppose K is not independent and let $\alpha \in K$ be a non-independent simplex. By definition, α has a face β such that $\bigcap \beta - \bigcup(\alpha - \beta) = \emptyset$ or, equivalently, $\bigcap \beta \subseteq \bigcup(\alpha - \beta)$. Therefore,

$$\begin{aligned} \mathbf{1}_{\bigcap \beta} &= \mathbf{1}_{\bigcap \beta} \cdot \mathbf{1}_{\bigcup(\alpha - \beta)} \\ &= \mathbf{1}_{\bigcap \beta} \cdot \sum_{\emptyset \neq \gamma \subseteq \alpha - \beta} (-1)^{\dim \gamma} \mathbf{1}_{\bigcap \gamma} \\ &= \sum_{\beta \subsetneq \delta \subseteq \alpha} (-1)^{\dim \delta - \dim \beta - 1} \mathbf{1}_{\bigcap \delta}, \end{aligned}$$

where $\delta = \beta \cup \gamma$, and $\dim \delta = \dim \beta + \dim \gamma + 1$ because $\beta \cap \gamma = \emptyset$. We therefore get

$$\sum_{\beta \subsetneq \delta \subseteq \alpha} (-1)^{\dim \delta} \mathbf{1}_{\bigcap \delta} = 0, \quad (5)$$

which implies that the set of faces of α that are cofaces of β is redundant. In other words, the minimality of K implies its independence.

Second realizability and boundary. Recall that a simplex β belongs to the boundary complex of the dual complex of B iff there is a point $y \in l(\beta)$ on the boundary of $\bigcup B$. By (ii) of the Inside-Outside Lemma, β also belongs to K . By (i) of the same lemma, every simplex in K for which there is no such point y is the face of a d -simplex. As explained after the proof of that lemma, every such $(d - 1)$ -simplex belongs to two d -simplices, one on each side. Intersect the (canonical images of the) simplices with an oriented line that avoids all simplices of dimension $d - 2$ or less. It meets the boundary $(d - 1)$ -simplices in some order, alternating between entering and exiting the underlying space. After entering and before exiting, the line may encounter a sequence of interior $(d - 1)$ -simplices, alternating between entering and exiting a d -simplex. Since this is true for almost all oriented lines, the mapping of abstract simplices to their canonical images is a geometric realization of K . Furthermore, the boundary complex and the underlying space of K are equal to those of the dual complex. This completes the proof of Theorem B.

5. CONCLUSION

The main result of this paper is a characterization of the minimal inclusion-exclusion formulas of a union of closed balls B in \mathbb{R}^d that correspond to simplicial complexes. What about inclusion-exclusion formulas that correspond to sets of simplices that do not form complexes? The central concept is that of an independent set of balls in \mathbb{R}^d , and our results rest on the observation that the maximum size of such a set is $d + 1$. There are other classes of geometric shapes with bounds on the size of independent sets. For example, the number of independent ovals (each bounded by an ellipse in \mathbb{R}^2) is at most five. Does an upper bound of $k + 1$ on the maximum number of independent shapes imply the existence of an abstract simplicial complex of dimension at most k that gives a correct inclusion-exclusion formula? The argument leading to Equation (5) might help in constructing such a complex. Can Theorems A and B be extended to ovals and other classes of simple shapes?

6. REFERENCES

- [1] F. AURENHAMMER. Voronoi diagrams—a survey of a fundamental geometric data structure. *ACM Comput. Surv.* **23** (1991), 345–405.
- [2] H. EDELSBRUNNER. The union of balls and its dual shape. *Discrete Comput. Geom.* **13** (1995), 415–440.
- [3] H. EDELSBRUNNER AND P. KOEHL. The geometry of biomolecular solvation. *Mathematical Sciences Research Institute Publications*, J. E. Goodman, J. Pach and E. Welzl (eds.), Cambridge Univ. Press, England, 241–273, 2005.
- [4] I. GELFAND, M. KAPRANOV AND A. ZELEVINSKY. *Discriminants, Resultants and Multidimensional Determinants*. Birkhäuser, Boston, 1994.
- [5] P. KOEHL. ProShape: understanding the shape of protein structures. Software at biogeometry.duke.edu/software/proshape, 2004.
- [6] K. KRATKY. The area of intersection of n equal circular disks. *J. Phys. A: Math. Gen.* **11** (1978), 1017–1024.
- [7] D. NAIMAN AND H. WYNN. Inclusion-exclusion Bonferroni identities and inequalities for discrete tube-like problems via Euler characteristics. *Ann. Statist.* **20** (1992), 43–76.
- [8] G. PERROT, B. CHENG, K. D. GIBSON, J. VILA, A. PALMER, A. NAYEEM, B. MAIGRET AND H. A. SCHERAGA. MSEED: a program for rapid determination of accessible surface areas and their derivatives. *J. Comput. Chem.* **13** (1992), 1–11.